This Alert Card contains important safety information that you need to be aware of before and during treatment with Remicade.

Show this card to any doctor involved in your treatment.

Please read the Remicade 'Package Leaflet' carefully before you start using this medicine.

When starting a new card, please keep this card as a reference for 4 months after this date.

REMICADE INTRAVENOUS INFUSION

Date aiven

DD	MM	YYYY
		YYYY
	MM	YYYY
		YYYY
	MM	YYYY

Next infusion due

MM	
MM	YYYY
MM	
MM	YYYY

Therapy initiation		DD / MM / YYY
Body weight (kg)	Dosage (mg/kg)	Dose administered (mg)

Please make sure you also have a list of all other medicines that you are using with you at any visit to a healthcare professional.

Keep this card with you for 4 months after your last dose of Remicade. Side effects may occur a long time after your last dose.



PATIENT ALERT CARD



Name Date of birth **Doctor's name** Doctor's telephone

Ask your doctor to record the typ for tuberculosis (TB) below:

Test	
Date	
Result	
Test	
Date	
Result	
Test	
Date	
Result	

Date of last screening(s)

Test
Date
Result
List of allergies
List of other medicines

IMPORTANT SAFETY INFORMATION

Infections

Before treatment with Remicade

- Tell your doctor if you have an infection even if it is a very minor one
- It is very important that you tell your doctor if you have ever had TB, or if you have been in close contact with someone who has had TB. Your doctor will test you to see if you have TB. Ask your doctor to record the type and date of your last screening(s) for TB on the card

Tell your doctor if you have hepatitis B or if you know or suspect you are a carrier of the hepatitis B virus

During treatment with Remicade

 Tell your doctor straight away if you have signs of an infection. Signs include a fever, feeling tired, (persistent) cough, shortness of breath, weight loss, night sweats, diarrhoea, wounds, dental problems, burning when urinating or 'flu-like' signs

Heart failure

Before treatment with Remicade

 Tell your doctor if you have any heart problems such as mild heart failure

During treatment with Remicade

 Tell your doctor straight away if you notice signs of a heart problem. Signs include shortness of breath, swelling of the feet or changes in your heartbeat Any suspected adverse events should be reported to the national spontaneous reporting system according to the national regulations.

SFDA (National pharmacovigilance and drug safety Center) Email to: npc.drug@sfda.gov.sa

Fax: +966-11-2057662 Online: http://ade.sfda.gov.sa/

Toll free number: 8002490000

Or you can contact company scientific office at: Email: GCC-PV2@its.ini.com

Fax: +966-11-2153190

