

PATIENT ALERT CARD

Your guide to safe beginning therapy
with JARDIANCE® empagliflozin

Keep this card with you all the times




Boehringer
Ingelheim

PO BOX 68423
Riyadh 11527
KSA

Jardiance®
(empagliflozin)



Make JARDIANCE part of your daily routine



Once-daily oral tablet



Take any time of day



With or without food

Tablet is not actual size.

Make sure to take JARDIANCE every day, even if you're feeling fine. This is important because skipping doses may cause your blood sugar level to increase again.

If you miss a dose, take it as soon as you remember. But, if a full day has passed, don't take 2 tablets—just continue with your normal dosage. Never take 2 tablets in a 24-hour period.



Try taking JARDIANCE at the same time every day, such as with breakfast or after you brush your teeth. This can help you to quickly make JARDIANCE part of your daily routine.

Follow your doctor's instructions when taking JARDIANCE®

- JARDIANCE® is an oral treatment for type 2 diabetes mellitus.
- JARDIANCE® is not approved for people with type 1 diabetes.
- JARDIANCE® is used to treat type 2 diabetes in adult patients (aged 18 years and older).
- JARDIANCE® is used if your diabetes cannot be controlled by diet and exercise alone and with other medicines for diabetes.
- JARDIANCE® is not approved for people with diabetic ketoacidosis (increased ketones in the blood or urine).
- JARDIANCE® may cause side effects, including:
Ketoacidosis is a rare serious life threatening condition due to severe reduction in the insulin which lead to increase ketones in your blood or urine.

Possible side effects with JARDIANCE


Like all medicines, JARDIANCE may cause side effects, although not everybody gets them. Carefully follow your doctor's advice if you experience any side effects.



Ketoacidosis has happened in people who have type 1 diabetes or type 2 diabetes, but less frequently in type 2 diabetes. Ketoacidosis and may need to be treated in a hospital. Ketoacidosis can happen even if blood glucose is not too elevated.

Diabetic Ketoacidosis Symptoms

- Loss of appetite, Stomach-area (abdominal) pain and vomiting (throwing up for more than 2 hours).
- Your breath smells fruity (acetone).
- You're tired, confused, or woozy.
- Thirst or a very dry mouth and skin.
- Feeling thirsty and urinating a lot.
- Rapid, deep breathing.



If you get any of the previous symptoms call your doctor or go to the emergency room right away.

JARDIANCE FAQs

Q How does JARDIANCE lower my blood sugar?

A JARDIANCE helps remove some of the extra sugar in your blood by passing it in your urine.

Q How do I take JARDIANCE?

A JARDIANCE is an oral tablet that you take once a day, with or without food.

Make the most of your doctor visits.

Call your doctor about any side effects

- If you experience any side effects, don't wait until your next visit to ask—call right away



If any questions come up before your next visit, try writing them down.



Here is a question that you may want to ask your doctor after starting JARDIANCE:

“Has my blood sugar (HbA1c) gone down since I started taking JARDIANCE?”

Any adverse reactions should be reported in accordance with the Saudi Vigilance spontaneous reporting system to:

National Pharmacovigilance and Drug Safety Centre NPC

Website <https://ade.sfda.gov.sa>

Email: npc.drug@sfda.gov.sa

Fax: +966-11-2057662

Call Center : 19999

Pharmacovigilance department in Boehringer Ingelheim:

Email: PV_local_Saudi_Arabia@boehringer-ingelheim.com

Phone:+966-11- 5116-500

This material approved by the Saudi food & drug authority

For extra copies please send email with the required amount & contact number to the below email:

PV_local_Saudi_Arabia@boehringer-ingelheim.com

V.2, Feb 2020

KSA/JAR-161194



**Boehringer
Ingelheim**

PO BOX 68423
Riyadh 11527
KSA

Jardiance® 
(empagliflozin)