COVER OUTSIDE - FRONT

Uptravi

film-coated tablets selexipag

Titration Guide

Starting Treatment With Uptravi

Please read the accompanying patient information leaflet before starting treatment.

Tell your doctor if you experience side effects, as your doctor may recommend that you change your Uptravi dose.

Tell your doctor if you are taking other medications as your doctor may recommend that you take Uptravi only once daily.

To reports any side effect(s):

The National Pharmacovigilance Centre (NPC):

SFDA Call Center: 19999 E-mail: npc.drug@sfda.gov.sa

Website: https://ade.sfda.gov.sa

If you have further question or require additional information, please contact our local safety department at:

Email: GCC-PV2@its.jnj.com

This document has been reviewed and approved by The Saudi Food and Drug Authority (SFDA).







MU_000236.pdf - Page 2 of 41 - November 19, 2021 - 05:26:48



Contents

How should you take Uptravi?	4
How should you step up your dose?	
What are the steps?	8
When should you step down?	Ю
Stenning down	12









When you move to your maintenance dose 14
If you forget to take Uptravi16
If you stop taking Uptravi
Titration diary









How should you take Uptravi?

Uptravi is a medicine taken every morning and evening for the treatment of pulmonary arterial hypertension, also called PAH.

The starting dose for Uptravi is 200 micrograms once in the morning and once in the evening.

The first intake of Uptravi should be in the evening.

You should take each dose with a glass of water, preferably during a meal.









There are 2 phases of treatment with Uptravi:

Titration

In the first several weeks, you will work with your doctor to find the dose of Uptravi that is right for you. Your doctor may have you step up from the starting dose to higher doses of Uptravi. Your doctor may step you down to a lower dose. This process is called titration. It lets your body gradually adjust to the medicine.

Maintenance

Once your doctor has found the dose that is right for you, this will be the dose you take on a regular basis. This is called the maintenance dose.











How should you step up your dose?

You will start at the 200 microgram dose in the morning and in the evening and after discussing with your doctor or nurse step up to the next dose.

The first intake of the increased dose should be in the evening. Each step usually lasts about 1 week. It could take several weeks to find the dose that is right for you.

The goal is to reach the dose that is most appropriate to treat you.

This dose will be your maintenance dose.









Every patient with PAH is different. **Not everyone will end up on the same maintenance dose.**

Some patients may have 200 micrograms in the morning and in the evening as their maintenance dose, while some will reach the highest dose of 1,600 micrograms in the morning and in the evening.

Others may reach a maintenance dose somewhere inbetween. What is important is that you reach the dose that is most appropriate to treat you.











The What are the steps?



Each dosing step lasts about 1 week.

STARTING DOSE



STEP 1

Morning: One 200 microgram tablet Evening: One 200 microgram tablet (total daily dose: 400 microgram)

STEP 2

Morning: Two 200 microgram tablets

Evening: Two 200 microgram tablets

(total daily dose: 800 microgram)



Morning: Three 200 microgram tablets

Evening: Three 200 microgram tablets (total daily dose: 1.200 microgram)

STEP 4

Morning: Four 200 microgram tablets **Evening:** Four 200 microgram tablets (total daily dose:

1,600 microgram)







800 microgram tablet (Use with steps 5 through 8 to reduce the number of tablets needed per dose.)



200 microgram tablet

HIGHEST 9 DOSE









STEP 5

Morning: One 800 and one 200 microgram tablet Evening: One 800 and one 200 microgram tablet (total daily dose: 2,000 microgram)

STEP 6

Morning: One 800 and two 200 microgram tablets Evening: One 800 and two 200 microgram tablets (total daily dose: 2.400 microgram)

STEP 7

Morning: One 800 and three 200 microgram tablets Evening: One 800 and three 200 microgram tablets (total daily dose: 2.800 microgram)

STEP 8

Morning: One 800 and four 200 microgram tablets Evening: One 800 and four 200 microgram tablets

(total daily dose:

3.200 microgram)

(Tablets are not actual size.)







When should you step down?

As with all medicines, you may experience side effects with Uptravi as you step up to higher doses.

Talk to your doctor or nurse if you have side effects. There are treatments available that can help relieve them.


The most common side effects (may affect more than 1 in 10 people) you may experience while taking Uptravi are:

- Headache Diarrhoea Nausea Vomiting Jaw pain Muscle pain
- Leg pain Joint pain Facial redness

For a full list of side effects see package leaflet for further information.









If you cannot tolerate the side effects even after your doctor or nurse has tried to treat them, he or she may recommend you step down to a lower dose.

If your doctor or nurse tells you to step down to a lower dose, you should take one less 200 microgram tablet in the morning and one less in the evening.

You should only step down after speaking with your PAH doctor or nurse. This stepping-down process will help you find the dose that is right for you, also called your maintenance dose.









Stepping down

START STEP 1

STEP 2

STEP 3

Each dosing step lasts about 1 week.



200 microgram tablet



Morning:
One 200 microgram tablet
Evening:

One 200 microgram tablet (total daily dose: 400 microgram) dose about every week by adding a 200 microgram tablet

to your morning and evening dose

(Tablets are not actual size.)









200 microgram tablet



800 microgram tablet

(Use with steps 5 through 8 to reduce the number of tablets needed per dose.)

 $MU_000236.pdf$ - Page 13 of 41 - November 19, 2021 - 05:26:48





When you move to your maintenance dose

The highest dose that you can tolerate during titration will become your **maintenance dose**. Your maintenance dose is the dose you should continue to take on a regular basis. Your doctor or nurse can prescribe an equivalent **single-tablet strength** for your maintenance dose.

This lets you take just one tablet in the morning and one in the evening, instead of multiple tablets for each dose.










For example, if your highest tolerated dose during titration was 1,200 micrograms once in the morning and once in the evening:



Over time, your doctor or nurse may adjust your maintenance dose as needed.











If you forget to take Uptravi

If you miss a dose, take the dose as soon as you remember, then continue to take your tablets at the usual times. If it is within 6 hours of when you would normally take your next dose, you should skip the missed dose and continue to take your medicine at the usual time.

Do not take a double dose to make up for a forgotten tablet.










If you stop taking Uptravi

Do not stop taking Uptravi unless your doctor or nurse tells you to. If, for any reason, you stop taking Uptravi for more than 3 consecutive days (if you missed 6 doses in a row), contact your PAH doctor or nurse immediately as your dose may need to be adjusted to avoid side effects.

Your doctor or nurse may have you resume treatment at a lower dose, gradually increasing to your previous maintenance dose.









Titration diary


Please read the instructions in the package leaflet carefully.

The following diary pages help you keep track of the number of tablets you need to take in the morning and evening during titration.

Use them to write down the number of tablets you take in the morning and the evening.

Each step usually lasts about 1 week, unless your doctor or nurse instructs you otherwise. If your titration steps last longer than 1 week there are additional diary pages to track this.

Use pages 20 to 27 to track the first weeks of treatment, when you are using 200 microgram tablets only (steps 1-4).












Write down your doctor or nurse's instructions:
Doctor's office telephone and e-mail:
Pharmacist's telephone:
Notes:








Every day write down in the boxes below how many tablets you take in the

20

morning and evening. I spoke with my doctor or nurse on DD/MM/YY.








Write down the number of the week of the treatment in the upper left hand corner. Every day write down in the boxes below how many tablets you take in the morning and evening. I spoke with my doctor or nurse on DD/MM/YY.

Date:





The first intake of an dose of Uptravi should be in the evening




Write down the number of the week of the treatment in the upper left hand corner. Every day write down in the boxes below how many tablets you take in the morning and evening. I spoke with my doctor or nurse on DD/MM/YY.















Write down the number of the week of the treatment in the upper left hand corner. Every day write down in the boxes below how many tablets you take in the morning and evening. I spoke with my doctor or nurse on DD/MM/YY.

Date:











Write down the number of the week of the treatment in the upper left hand corner. Every day write down in the boxes below how many tablets you take in the morning and evening. I spoke with my doctor or nurse on DD/MM/YY.

Date:








Write down the number of the week of the treatment in the upper left hand corner. Every day write down in the boxes below how many tablets you take in the morning and evening. I spoke with my doctor or nurse on DD/MM/YY.

Date:













Write down the number of the week of the treatment in the upper left hand corner. Every day write down in the boxes below how many tablets you take in the morning and evening. I spoke with my doctor or nurse on DD/MM/YY.

Date:







Write down the number of the week of the treatment in the upper left hand corner. Every day write down in the boxes below how many tablets you take in the morning and evening. I spoke with my doctor or nurse on DD/MM/YY.

Date:



MU 000236.pdf - Page 27 of 41 - November 19, 2021 - 05:26:48







200 microgram tablet



800 microgram tablet (Use with steps 5 through 8 to reduce the number of tablets needed per dose.)

Use the following diary pages if your doctor or nurse prescribes 800 microgram tablets in addition to your 200 microgram tablets.

On the diary pages, check off that you have taken **one** 800 microgram tablet every day in the morning and in the evening with your prescribed number of 200 microgram tablets.









Remember to talk to your PAH doctor or nurse regularly.

Notes:

Write down your doctor or nurse's instructions:
Doctor's office telephone and e-mail:
Pharmacist's telephone:






Write down the number of the week of the treatment in the upper left hand corner. Every day write down in the boxes below how many tablets you take in the morning and evening. I spoke with my doctor or nurse on DD/MM/YY.

30








Write down the number of the week of the treatment in the upper left hand corner. Every day write down in the boxes below how many tablets you take in the morning and evening. I spoke with my doctor or nurse on DD/MM/YY.










Write down the number of the week of the treatment in the upper left hand corner. Every day write down in the boxes below how many tablets you take in the morning and evening. I spoke with my doctor or nurse on DD/MM/YY.










Write down the number of the week of the treatment in the upper left hand corner. Every day write down in the boxes below how many tablets you take in the morning and evening. I spoke with my doctor or nurse on DD/MM/YY.








Write down the number of the week of the treatment in the upper left hand corner. Every day write down in the boxes below how many tablets you take in the morning and evening. I spoke with my doctor or nurse on DD/MM/YY.










Write down the number of the week of the treatment in the upper left hand corner. Every day write down in the boxes below how many tablets you take in the morning and evening. I spoke with my doctor or nurse on DD/MM/YY.








Write down the number of the week of the treatment in the upper left hand corner. Every day write down in the boxes below how many tablets you take in the morning and evening. I spoke with my doctor or nurse on DD/MM/YY.












Write down the number of the week of the treatment in the upper left hand corner. Every day write down in the boxes below how many tablets you take in the morning and evening. I spoke with my doctor or nurse on DD/MM/YY.










Notes






















Product	Product Artwork CoE	S		Janssen 🖊
Shared ma	ilbox: RA-JAN	VBE-GR/	Shared mailbox: RA-JANBE-GRAPHDES@its.jnj.com	
Booklet		UPTRAVI	M	
PC	M	00	MU_000236	
Template f	Template Ref. Name:	Allpack 120x65	Allpack/Booklet/120x65/V1 120x65	
Tech. info/Spec:	Spec:	,		
Market:		SA		
Mat. ID Code:	de:	N/A		
File Name:	٠.	MU_00	MU_000236.indd (CC - PC)	
Replaces PCC:	PCC:	N/A		
Font Type:		Gotham	E	
Font Size:		8 pt		
x-height	Times Nev	v Romar	Times New Roman compared to Actelion standard fonts	n standard fonts
(lowercase) Times NR	Times New Roman	Roman	Helvetica Neue LT / Arial	Univers















Univers 5.4 pt xxx 7.2 pt xxx 6.3 pt xxx 8 pt

Arial 6.1 pt 7 pt

> 8 pt xxx XXX 9 pt

(Designer) 1.03-NOV-21 (Syed) 2.05-NOV-21 (Syed) 3.19-NOV-21 (Syed)

XXX

XXX 7.8 pt xxx

7 pt xxx

1.104 mm 1,262 mm 1,42 mm Black Cyan

0,947 mm

Magenta Yellow



























































