

Patient Card

Methotrexate

2.5 mg Tablet

Methotrexate

SANDOZ

Patient card for Methotrexate: Medication errors

Oral formulations of methotrexate-containing medicinal products with at least one indication requiring treatment once a week:

This patient card is only intended for patients who use a methotrexate-containing medicine for indications inserted by the marketing authorization holders (mahs) in grouped terms, e.g. colitis, arthritis, psoriasis, as per indications listed in package leaflet.

If you use methotrexate for one of the above mentioned indications, you should only take methotrexate once a week

Write here in full the day of the week for intake: _____

Do not take more than the prescribed dose.

Overdose could lead to serious adverse effects and may be fatal. Symptoms of overdose are e.g. sore throat, fever, mouth ulcers, diarrhea, vomiting, skin rashes, bleeding or unusual weakness.

If you think you have taken more than the prescribed dose, consult a physician immediately.

Always show this card to health care professionals not familiar with your methotrexate treatment to alert them about your once weekly use (e.g. on hospital admission, change of care).

For more information, please read the patient leaflet inserted in the package.

This document has been approved by Saudi Food and Drug Authority (SFDA).



Sandoz reporting adverse events:
Email: adverse.event.sau@sandoz.com
Website: <https://pvi1j.solutions.iqvia.com/pvi-web/>



Saudi Food and Drug Authority National Pharmacovigilance Center
Unified Contact Center: 19999
Email: npc.drug@sfda.gov.sa
Website: <https://ade.sfda.gov.sa>

SANDOZ