



## Patient Reminder Card

This document has been reviewed and approved by the Saudi Food and Drug Authority (SFDA).

**IXIFI 100 mg**  
**Infiximab**

Show this card to any physician involved in your treatment.

This Patient Reminder Card contains important safety information that you need to be aware of before and during treatment with IXIFI.<sup>1</sup>

**Patient Name:** \_\_\_\_\_ **Physician Name:** \_\_\_\_\_ **Telephone no.:** \_\_\_\_\_

It is important that you and your physician record the brand name and batch number of your medication.

Date of IXIFI therapy initiation: \_\_\_\_\_ Dates of administration in the current cycle: \_\_\_\_\_

- When initiating a new card, ensure you retain this card for reference for four months following your last dose of IXIFI.
- Read the IXIFI 'Patient Leaflet' carefully before you start using this medicine.
- Keep this card with you for 4 months after your last dose of IXIFI, or in case of pregnancy, for 12 months after the birth of your baby.
- Side effects may occur after a prolonged use of the drug.

Ask your physician to record the type and date of the last tuberculosis (TB) screening test below:

Test: \_\_\_\_\_ \ \_\_\_\_\_ Date: \_\_\_\_\_ \ \_\_\_\_\_ Result: \_\_\_\_\_ \ \_\_\_\_\_

List of known sensitivities (allergies): \_\_\_\_\_ List of other medicines you are taking: \_\_\_\_\_

### **Infections**

#### **Before using IXIFI**

- Tell your physician if you have any infection or get a lot of infections or have infections that keep coming back.<sup>1</sup>
- It is very important that you tell your physician if you have ever had tuberculosis (TB), or if you have been in close contact with someone with TB.<sup>1</sup>  
Your physician will test you to see if you have TB.<sup>1</sup> Your doctor should monitor you closely for signs and symptoms of TB during treatment with IXIFI.<sup>1</sup>
- Tell your physician if you have hepatitis B or if you know or suspect that you are a carrier of the virus that causes this disease.<sup>1</sup>
- Tell your physician if you have signs of an infection, such as a fever, cough, flu-like symptoms.<sup>1</sup>
- Tell your physician if you have any open cuts or sores on your body.<sup>1</sup>
- Tell your physician if you have diabetes or an immune system problem. People with these conditions have a higher chance for infections.<sup>1</sup>
- Tell your physician if you live or have lived in certain parts of the country (such as the Ohio and Mississippi River valleys) where there is an increased risk for getting certain kinds of fungal infections (histoplasmosis, coccidioidomycosis, or blastomycosis). These infections may develop or become more severe if you receive IXIFI. If you do not know if you have lived in an area where histoplasmosis, coccidioidomycosis, or blastomycosis is common, ask your doctor.<sup>1</sup>



### During Treatment with IXIFI

- Call your doctor right away if you have an infection, any sign of an infection including a fever, cough, flu-like symptoms, or have open cuts or sores on your body, call your doctor right away.<sup>1</sup>

### Heart Problems

#### Before using IXIFI

- Tell your doctor if you have heart failure or other heart conditions.<sup>1</sup>

### During Treatment with IXIFI

- Tell your doctor of any new or worse symptoms of heart failure including shortness of breath, swelling of ankles or feet, or sudden weight gain.<sup>1</sup>

### Pregnancy and Vaccinations

If you received IXIFI while you were pregnant, your baby may have a higher risk for getting an infection.<sup>1</sup>

If you were receiving IXIFI during your pregnancy, it is crucial to inform your baby's doctor and other healthcare professionals about your IXIFI use. This will help them determine the appropriate timing for your baby's vaccinations.<sup>1</sup>

Your baby must not receive a "live vaccine" such as BCG (intended to prevent tuberculosis) within 12 months after birth.<sup>1,2</sup>

At any visit to a healthcare professional, please make sure to **tell your doctor about all the medicines you take**, including prescription and over-the-counter medicines, vitamins, and herbal supplements.<sup>1</sup>

Keep this card with you for 4 months after receiving the last dose of IXIFI.<sup>1</sup>

### Reporting of Suspected Adverse Reactions:

Adverse events can be reported directly to:

#### The National Pharmacovigilance Center (NPC) at Saudi Food and Drug Authority (SFDA)

SFDA call center: 19999

E-mail: [npc.drug@sfd.gov.sa](mailto:npc.drug@sfd.gov.sa)

Website: <http://ade.sfd.gov.sa>

### Pharmacovigilance Department in the Company

E-mail: [SAU.AEReporting@pfizer.com](mailto:SAU.AEReporting@pfizer.com)

For extra copies, please send an email with your contact details and the required amount to

[SAU.AEReporting@pfizer.com](mailto:SAU.AEReporting@pfizer.com)

#### References:

1. Ixifi Package Leaflet. Saudi Arabia, December 2023.
2. Ixifi LPD. Saudi Arabia, December 2023.



Pfizer Scientific Technical Limited Company:  
Al Riyadh – Al Aqeeq Dis.  
King Abdullah Financial District, Building 4.07  
Postal Code 13519  
Tel: +966 12 2293500  
Fax: +966 12 2293692

Version Number: 1  
Date of Approval: December - 2024.

PP-IFX-SAU-0033