

Important Risk Minimization Information for Patients

Reminder Card

Feriprox Modified Release Tablets (Deferiprone) 1000mg

Check twice

Deferiprone 1000 mg tablets are taken
two-times-a-day

NEW



Deferiprone 500 mg tablets are taken
three-times-a-day



Deferiprone 100 mg/mL oral solution is taken
three-times-a-day

Check twice

to ensure you are taking
your medication as
prescribed by your doctor.

Review your dosing
schedule carefully to avoid
any medication errors.

Please report any suspected adverse reactions to
The National Pharmacovigilance Centre (NPC), SFDA:

- SFDA Call Center: 19999
- E-mail: npc.drug@sfd.gov.sa
- Website: <https://ade.sfd.gov.sa>

Or

LOCAL REPRESENTATIVE OF THE MARKETING AUTHORISATION HOLDER IN KSA
Biologix FZCO, Hibatullah Al Ghaffari Street- Suleimaniah
Kingdom of Saudi Arabia P.O.Box 991, Riyadh 11421 Biologix Pharmacovigilance Department:
• E-mail: Pharmacovigilance-ksa@biologixpharma.com
• Tel + 966 11 4646 955 Ext 286
• Mobile: +966559994037

Ferriprox is available in three formulations:

- 1000 mg Modified Release tablets two-times-a-day
- 500 mg tablets three-times-a-day
- 100 mg/mL solution three-times-a-day

**Be sure to review
your medication
formulation, dose, and
dosing schedule with
your doctor carefully
and make note
of any changes.**



What is Ferriprox and Ferriprox MR?

- Ferriprox contains the active substance deferiprone. Ferriprox is an iron chelator, a type of medicine that is used to remove excess iron from the body.
- Ferriprox is used for the treatment of iron overload in patients with thalassaemia major when current chelation therapy is inadequate.
- Ferriprox can be used in combination with another iron chelator if your doctor has determined that treatment with any iron chelator alone has been ineffective or if your condition requires rapid or intensive correction of the excess iron in your body.

For more information, please refer to the patient information leaflet inside Ferriprox pack.