Patient Reminder Card for Capitan® 10 & 20mg (Isotretinoin)

You have been given this card because you have been prescribed Capitan® 10 & 20mg (Isotretinoin) and you need to know the possible side effects.

Please read this card and the patient information leaflet thoroughly before starting your treatment with Capitan® 10 & 20mg (Isotretinoin) to ensure that you understand the range of side effects associated with taking isotretinoin.

Please refer to the Patient Information Leaflet (PIL) for Capitan® 10 & 20mg (Isotretinoin) for more information.

Pregnancy risks

- YOU MUST NOT take Isotretinoin during pregnancy or if you are planning to get pregnant.
- Exposure to Isotretinoin can seriously harm an unborn baby.
- You must avoid becoming pregnant in the month before starting isotretinoin treatment,
 while taking isotretinoin, and for 1 month after stopping treatment.
- If you may be able to get pregnant, you must use effective contraception during this time.

 Talk to your prescriber to make sure you are using the right kind of contraception.
- If you miss your period, become pregnant, or think you might be pregnant, stop taking isotretinoin immediately, contact your dermatologist, and seek medical advice. This applies during isotretinoin treatment and for 1 month after stopping.
- If there is any chance you could get pregnant, you must have regular pregnancy testing as instructed by your dermatologist. For further information, refer to SPC or PIL.



Important Reminders for Patients

- **DO NOT** share this medicine with anybody. Any Capitan® 10 & 20mg soft capsules unused after the end of treatment or by the expiry date must be returned in their original packaging to the place of supply (physician or pharmacist) for proper disposal.
- You MUST attend clinic appointments during treatment for regular monitoring and care.
- Mental health: Talk to your dermatologist about your treatment or if you have any concerns.
- You should stop taking Capitan® 10 & 20mg (Isotretinoin) and contact your dermatologist for further advice if you have serious concerns about your mental health or thoughts of harming yourself or other serious side effects.
- You must NOT donate blood during treatment with Capitan® 10 & 20mg (Isotretinoin) and for 1 month after stopping.

If you have any questions or concerns about taking Capitan® 10 & 20mg 10 & 20mg (Isotretinoin), talk to your dermatologist.

Call For Reporting:

Any suspected adverse reactions should be reported immediately to Avalon Pharma Saudi or to the National Pharmacovigilance Center (NPC) at Saudi food and drug authority.

Pharmacovigilance Department at Middle East Pharmaceutical Industries Company (Avalon Pharma):

• Email: pharmacovigilance@avalon.com.sa

Website: https://www.avalonpharmaceutical.com/

Phone No: +966 55 222 1704

The National Pharmacovigilance Center (NPC): (Saudi food and drug authority)

Email: npc.drug@sfda.gov.sa

Call Center: 19999

Website: https://ade.sfda.gov.sa/

• QR Code:



